

DON'T LET LEAD HITCH A RIDE HOME



Breathing or swallowing lead dust can cause




Brain damage Memory loss Trouble concentrating
Depression Nerve damage Learning disabilities
Headaches Trouble sleeping Aggressive behavior
Irritability Lower IQ



High blood pressure



Abnormal sperm
Impotence
Loss of sex drive
Infertility
Miscarriage, preterm labor, or stillbirth



Muscle or joint pain

Constipation
Kidney damage
Loss of appetite
Stomach cramps

Most kids and adults with lead poisoning
do **NOT** feel or look sick.

It can take years for
the effects of lead exposure to appear.

Many jobs and hobbies may expose you to lead including:
plastic/glass manufacturing, automobile repair, radiator
repair, battery manufacturing, iron/steel working,
welding, soldering, mining, road construction,
firing ranges, demolition, renovation

PROTECT YOUR FAMILY FROM LEAD

Keep yourself and your family safe:



Wash your hands before
eating, drinking, smoking or
touching anything



Change your clothes and shoes
before going home or getting
into your car



Wash lead-exposed clothes
separately from regular laundry



Use a HEPA vacuum to clean
lead dust from your car and
home



Talk to your doctor about a lead
test if you suspect exposure

To learn more visit nvclppp.org
or call 702-895-5067



This publication was supported by the Grant or Cooperative Agreement Number, 1 NU2E2 H001366-01-00, funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services.